

Scurlock Tower 6560 Fannin, Suite 1750 Houston, TX 77030 713-790-0400

Nuclear Stress Test Instructions

Food:	No food or drink for 3 hours before the test (except water)Please bring a light snack and a bottle of water	
Caffeine	 No caffeinated or decaffeinated products 24 hours prior This includes soft drinks, tea, coffee, chocolate, energy drinks, excedrin, anacin 	
Clothing	Wear comfortable clothingIf exercising, wear running shoes and bring a small towel	
Notes	 Test time will range 3/12 to 4 hours If pregnant or planning pregnancy, let us know prior to the test For weight over 325lbs, the test will be done over 2 days 	
Medications	• Take all your medication prior to the test EXCEPT:	

e e	In (non treadmill) test owing 24 hours prior:	If you are doing the treadmill test
Aggrenox	Dipyridamole	<u>Do not take Beta blockers</u> for 48 hours prior:
		Metoprolol
Theo-Dur	Aminophylline	Carvedilol
		Bisoprolol
Persantine	Theophylline	Atenolol
	1 2	Bystolic
**If doing the Lexiscan, pl	ease bring a caffeinated	Propranolol
beverage of your choice to		Labetalol
caffeine will reverse the eff		Sotalol
quickly. (<u>Do not drink</u> it p		Nebivolol

Cancellations • Cancellations with less than 24h notice will be subject to a cancellation fee of \$300

- You must contact us during business hours. After-hours cancellations may not be honored.
- The \$300 cancellation fee covers the nuclear material we have ordered in preparation for your test.